


# Leptospirosis in Dairy Cattle

Leptospirosis is a bacterial infection and is an important zoonotic disease in New Zealand especially for people working with livestock such as farmers, abattoir workers and veterinarians.

The bacteria colonises the kidney and genital tract of a maintenance host (cattle, sheep, pigs, rats, possums and hedgehogs are all implicated here) and is shed in the urine which is a major source of contamination and infection to accidental hosts (sheep, deer and other animals as well as humans) in which clinical disease can occur.



Leptospire can survive in wet soil and stagnant water and the likelihood of infection increases with heavy rainfall and flooding. Leptospire do not replicate outside of the host but their numbers in the environment can be maintained by constant shedding by infected animals.

Transmission is through mucous membranes of the nose, eyes, mouth, reproductive tract, digestive tract or breaks in the skin. Leptospire can also be shed through aborted fetuses, placental fluids and milk.

There are several serovars but the most common in New Zealand are *L. hardjo-bovis*, *L. tarassovi*, *L. ballum*, and *L. pomona*. The maintenance hosts are:

Cattle: *L. hardjo-bovis*, *L. pomona*

Sheep: *L. hardjo-bovis*, *L. pomona*

Pigs: *L. pomona*, *L. tarassovi*

Rats: *L. tarassovi*, *L. ballum*

Dogs: *L. copenhageni*.

Infection will cause mild clinical signs and sporadic abortion in the maintenance hosts. Cattle that become infected with *L. pomona* can present with severe clinical disease.

Pigs are a major risk factor when they present on cattle farms. Ideally pigs should be vaccinated and their effluent kept far away from contact with cattle. Sufficient vaccination programmes are even more crucial on these farms.

## Zoonosis

There are approximately 140 reported cases per year of leptospirosis in humans and many more non reported cases. High risk occupations are farmers and farm workers, meat processors and veterinarians. Infection occurs through direct contact with urine or contaminated soil or water and through inhalation of aerosolised droplets.

There are increased rates of human infection with *L. Tarassovi* in recent years - this strain is not in vaccines.

Symptoms of leptospirosis can be confused with influenza. Fevers, chills, headache, nausea, cough, abdominal pain, muscle pain, and diarrhoea are

symptoms. In severe cases jaundice, kidney failure, meningitis, haemorrhage and hemodynamic collapse may occur. Many cases require hospitalisation and there can be long lasting symptoms. Clinical illness varies from mild to severe with several weeks off work. Studies have shown that about 20% of infected people develop clinical signs and about 6% suffer severe clinical disease.



## Signs of disease in cattle

### *L. hardjo-bovis* & *tarassovi*

Infected cattle may not show obvious clinical signs, sometimes they have a fever, or just appear unwell or have loss of appetite. Lactating cows may develop mastitis or a drop in production. Abortion or stillbirths can result even 6 to 12 weeks after infection. It was estimated that about 10% abortions in beef cattle may be caused by *L. hardjo-bovis* or *L. pomona*. Chronic infection of the genital tract may result in infertility, abortion, stillbirths, and birth of weak calves. Most commonly cattle show no clinical signs with *L. hardjo-bovis* or *L. tarassovi* infection but can create a carrier state in the kidneys associated with long term urinary shedding.

### *L. pomona*

Like *L. hardjo-bovis*, *L. pomona* can cause a fever and inappetence which may go unnoticed and can also cause abortion, still births and birth of weak calves. Abortion can occur after the 4th month of gestation during the acute stage or 6-12 weeks after infection. Most commonly, abortion occurs in the 6<sup>th</sup> or 7<sup>th</sup> month of gestation. Mastitis and a drop in production also occur. Mastitis due to leptospirosis does not respond to treatment and does not create an inflammatory reaction in the udder but does cause milk to resemble colostrum. Calves can suffer greatly from *L. pomona* infection; they will have a high fever, depression, haemolytic anaemia, jaundice, pale mucous membranes, and red water and may die within 3-5 days.



## Diagnosis

In cases of abortion there is rarely diagnostic lesions observed and it is very difficult to isolate leptospire for a definitive diagnosis but it can be attempted. Blood testing the cow and foetus for serovar-specific antibodies is the best diagnostic tool to use. Antibody titres of 1:100 and higher indicate that animals are likely to shed *Leptospira* in the urine.

A real time PCR is also commercially available for testing the presence of leptospire in urine from live animals or kidney tissue of an aborted foetus.

Post mortem examinations provide more clues in cases of acute infection with *Leptospira*. Infection can cause kidney and liver disease as well as red blood cells being destroyed which results in anaemia.

## Treatment

Animals can be treated with appropriate antibiotics prescribed by your veterinarian. The aim of treatment is to prevent irreparable damage to the kidneys and liver but often by the time the animal is presented it is too late for antibiotics to be effective. Treatment will eliminate shedding of leptospire.

## Herd health planning: Prevention and control

Farms are at risk if they have high rainfall during spring and autumn, experience flooding, and have waterways through the property and stagnant water pools in paddocks. Rodents (particularly rats), feral pigs, deer, sheep, possums, hedgehogs and rabbits may also be a source of infection. Livestock that are in contact with unvaccinated stock while grazing off farm are also susceptible

Vaccination is the corner stone of leptospire risk management plans. Other preventative measures include fencing off of water ways, excellent rodent control especially around feed storage and high standards of personal hygiene when working with cattle and pigs - covering cuts with waterproof plasters and no smoking, eating or drinking particular while in the milking shed.

## Vaccination

The currently available vaccines are very effective in protecting the animal from shedding leptospire in their urine but only for the serovars hardjo-bovis, pomona and copenhageni. **Note: *L. tarassovi* is not protected by the vaccines.**

Animals vaccinated after being exposed to infection can continue to shed as adults. Therefore the older a calf is at the time of first vaccination the greater the likelihood of having already been exposed to the bacteria.

Vaccines are efficient at preventing urine shedding (80-100% efficacy), if the animals are vaccinated before exposure to the bacteria. Recent research has shown that maternal derived antibody, which the calf receives from colostrum, is unlikely to interfere with vaccine efficacy. Most calves will have lost maternal protection by four to six weeks of age.

With dairy calves the recommended vaccination programme is to administer the first vaccinations ten weeks after the start of calving followed by a booster 4-6 weeks later. It is strongly recommended to have the vaccinations completed by the time the calves are 4 months old, i.e. Christmas.

At a minimum they should be completed by six months of age. A third vaccination should be given at the time of whole herd booster vaccination in the autumn

### SUMMARY:

- Effective and robust vaccination programme: all stock, early calf vaccination, correct booster intervals.
- Excellent rodent control especially around feed storage areas.
- High level of personal hygiene: cover cuts, avoid hand-to-mouth contact while working with the animals.
- Fence of waterways and effluent areas.

Information and illustration material for this fact sheet was kindly provided by Massey University and the Farmer Leptospirosis Action Group, a project of the Sustainable Farming Fund (SFF/MPI). Information available at: [www.leptospirosis.org.nz](http://www.leptospirosis.org.nz)

For more information contact your local XLVets practice:

**XLVets**  
Excellence in Practice

 **WELFARM**